

Energisation of interstellar media and cosmic ray production by jets from X-ray binaries

R. P. Fender^{1,2}, T.J. Maccarone², Z. van Kesteren²

¹*School of Physics and Astronomy, University of Southampton Hampshire SO17 1BJ*

²*Astronomical Institute ‘Anton Pannekoek’, University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands*

2 February 2008

ABSTRACT

Drawing on recent estimates of the power of jets from X-ray binary systems as a function of X-ray luminosity, combined with improved estimates of the relevant $\text{Log}(N)$ - $\text{Log}(L_X)$ luminosity functions, we calculate the total energy input to the interstellar medium (ISM) from these objects. The input of kinetic energy to the ISM via jets is dominated by those of the black hole systems, in contrast to the radiative input, which is dominated by accreting neutron stars. Summing the energy input from black hole jets L_J in the Milky Way, we find that it is likely to correspond to $\geq 1\%$ of L_{SNe} , the time-averaged kinetic luminosity of supernovae, and $\geq 5\%$ of L_{CR} , the cosmic ray luminosity. Given uncertainties in jet power estimates, significantly larger contributions are possible. Furthermore, in elliptical galaxies with comparable distributions of low mass X-ray binaries, but far fewer supernovae, the ratio L_J/L_{SNe} is likely to be larger by a factor of ~ 5 . We conclude that jets from X-ray binaries may be an important, distributed, source of kinetic energy to the ISM in the form of relativistic shocks, and as a result are likely to be a major source of cosmic rays.

Key words:

binaries: close – ISM:jets and outflows – black hole physics

1 INTRODUCTION

The interstellar medium (ISM) in the Milky Way and other galaxies receives a continuous input of energy in the form of both bulk motions and radiation. The result of this energy input is the heating of the ISM, the generation of interstellar turbulence and the production of high energy cosmic rays. The most violent processes are those associated with stellar evolution: stellar winds, protostellar jets and supernovae (SNe). More specifically, cosmic ray production is only likely to be associated with high velocity (i.e. $\gtrsim 0.1c$) shocks, and so their most likely source has long been considered to be the SNe. See e.g. Chevalier (1977), McCray & Snow (1979) and Elmegreen & Scalo (2004) for reviews of these processes.

However, in recent years it has become clear that there is at least one other potential source of powerful relativistic shocks in the ISM, namely the jets resulting from accretion onto stellar mass black holes and neutron stars in X-ray binary systems (Mirabel & Rodriguez 1999; Fender 2005). These jets seem to be powerful and ubiquitous in the majority of such systems which exist in ‘hard’ X-ray spectral states and/or are undergoing X-ray outbursts (Fender 2001; Fender, Belloni & Gallo 2004). Heinz & Sunyaev (2002) estimated the total power input from a handful of the most

powerful jets in our galaxy and concluded that they might contribute 1–10% of the cosmic ray luminosity of the galaxy, perhaps with a specific spectral signature. However, recent advances in our estimates of the scaling of jet power with X-ray luminosity (Fender, Gallo & Jonker 2003; Fender, Belloni & Gallo 2004), combined with improvements in our knowledge of the distribution of X-ray sources as a function of luminosity (e.g. Grimm, Gilfanov & Sunyaev 2002, 2003; Gilfanov 2004) have allowed us to recalculate more accurately the sum of the power input from such jets. In this paper we present the results of these calculations.

2 LUMINOSITY FUNCTIONS

2.1 Differential luminosity function for LMXBs

Grimm et al. (2003) presented the X-ray luminosity function (XLF) for LMXBs in the Milky Way in the form of a cut-off power law. Since then, Gilfanov (2004) has produced an average differential XLF for LMXBs in *all* galaxies. This function is in the form of a power law with two breaks:




Figure 1. Estimated total numbers of X-ray binaries in the Milky Way as a function of X-ray luminosity. At the highest luminosities, above $10^{35} \text{ erg s}^{-1}$, the low-mass X-ray binaries (LMXBs) dominate the source population; although at even lower luminosities numbers of high-mass X-ray binaries (HMXBs) may come to dominate, the numbers are not well measured at those levels. Since the process of disc-jet coupling does not seem to be strongly affected by the nature of the companion star (see e.g. discussion in Fender 2005) this figure illustrates that the total X-ray binary jet power into the interstellar medium will be dominated by the LMXBs.

$$\frac{dN}{dL_{X,38}} = \begin{cases} K_1 (L_{X,38}/L_{b,1})^{-\alpha_1} & L_{X,38} < L_{b,1} \\ K_2 (L_{X,38}/L_{b,2})^{-\alpha_2} & L_{b,1} < L_{X,38} < L_{b,2} \\ K_3 (L_{X,38}/L_{cut})^{-\alpha_3} & L_{b,2} < L_{X,38} < L_{cut} \\ 0 & L_{X,38} > L_{cut} \end{cases} \quad (1)$$

where $L_{X,38} = L_X/10^{38} \text{ erg s}^{-1}$ and normalizations $K_{1,2,3}$ are related by

$$K_2 = K_1 (L_{b,1}/L_{b,2})^{\alpha_2}$$

$$K_3 = K_2 (L_{b,2}/L_{cut})^{\alpha_3}$$

The best fit to the overall normalisation is given as

$$K_1 = 440.4 \pm 25.9 \text{ per } 10^{11} M_\odot \quad (2)$$

In the following we shall use this as the template LMXB XLF for all galaxies. Based on Liu et al. (2000, 2001) we estimate the black hole and neutron star fractions amongst the LMXBs to be $f_{BH} = 0.2$, $f_{NS} = 0.8$. This luminosity function, normalised to the Milky Way using a total stellar mass of $4.5 \times 10^{10} M_\odot$ (Gilfanov 2004) is plotted in Fig 1.

2.2 Differential luminosity function for HMXBs

For HMXBs, we use the differential form of the XLF from Grimm et al. (2003), which is simply a steep power law:

$$\frac{dN}{dL_{X,38}} = K L_{X,38}^{-1.64}, \quad (3)$$

where for the Milky Way $K \sim 0.7$. Based on Liu et al. (2000, 2001) we estimate the black hole and neutron star fractions amongst the HMXBs to be $f_{BH} = 0.05$, $f_{NS} = 0.95$. This function is plotted in Fig 1 alongside the LMXB XLF.

2.3 $L_J(L_X)$ for X-ray binary jets

2.3.1 Steady ('low/hard' state) jets

Corbel et al. (2000; 2003) and Gallo, Fender & Pooley (2003) have found an apparently universal correlation between radio and X-ray luminosities for black hole candidate (BHC) binaries in the 'low/hard' and 'quiescent' X-ray states (which seem to correspond to accretion rates below about 1–10% of the Eddington limit). The relation has the form

$$L_{\text{radio}} \propto L_X^b \quad (4)$$

where $b \sim 0.7$. The relation between jet and radio luminosities may also be considered to have a power-law form

$$L_{\text{radio}} \propto L_J^c \quad (5)$$

Several models of steady, conical jets predict $c \sim 1.4$ (e.g. Blandford & Königl 1979; Heinz & Sunyaev 2003), which leads to

$$L_J = A_{\text{steady,BHC}} L_X^{0.5} \quad (6)$$

in Eddington units, or

$$L_{J,38} = A_{\text{steady,BHC}} (L_{X,38})^{0.5} \left(\frac{M}{M_\odot} \right)^{0.5} \quad (7)$$

indicating that the fractional jet power increases rapidly as the X-ray luminosity decreases (Fender, Gallo & Jonker 2003; Fender, Belloni & Gallo 2004).

The value of the normalisation $A_{\text{steady,BHC}}$ is uncertain. Fender, Gallo & Jonker (2003) estimated (in their opinion conservatively) that $A_{\text{steady,BHC}} \geq 6 \times 10^{-3}$. While it is certainly far from the consensus, larger values of $A_{\text{steady,BHC}}$ may be more realistic. Malzac, Merloni & Fabian (2004) have suggested that even at $L_X \sim 10^{-3} L_{\text{Edd}}$, the jet may be an order of magnitude more powerful than the X-ray emission, which corresponds to $A_{\text{steady,BHC}} \sim 0.3$. Alternatively, we can set the value of $A_{\text{steady,BHC}}$ so that it corresponds to the transition from the low/hard to high/soft X-ray states (see e.g. McClintock & Remillard 2005 for a discussion), in which case $A_{\text{BHC}} \sim 0.1$. In all the following discussions we shall consider $0.006 \leq A_{\text{BHC}} \leq 0.3$ as covering the full range of likely values.

We further assume that $A_{\text{steady,NS}} = 0.1 A_{\text{steady,BHC}}$, based on the lower 'radio loudness' of NS XRBs (Fender & Kuulkers 2001; Migliari et al. 2003; Fender, Gallo & Jonker 2003; Munro et al. 2004; Migliari & Fender in prep). It should be stressed that the $L_{\text{radio}} \propto L_X^{0.7}$ relation has *not* been established for neutron star X-ray binaries, and is only assumed in Fender, Gallo & Jonker (2003) by analogy with the BHCs. However, since the BHCs are so much more radio loud then this is not significant for the total energy budget calculated here.

2.3.2 $L_J(L_X)$ for transient jets

A fit to the power in transient optically thin ejection events from *black hole* X-ray binaries as a function of X-ray luminosity (Fender, Belloni & Gallo 2004), simplified to one significant figure, gives

$$L_J = A_{\text{trans,BHC}} L_X^{0.5} \quad (8)$$

in Eddington units, where $A_{\text{trans,BHC}} \sim 0.4$. The same $L_X^{0.5}$ dependence (see Fender, Belloni & Gallo 2004 for caveats) as for the steady jets means the equation has the same form with simply a different normalisation:

$$L_{J,38} = A_{\text{trans,BHC}}(L_{X,38})^{0.5} \left(\frac{M}{M_\odot} \right)^{0.5} \quad (9)$$

It should be stressed that the normalisation is also rather uncertain, being based on a small sample of sources for which the energy estimate is itself intrinsically uncertain and different to that used for the steady jet sources (see discussion in Fender, Belloni & Gallo 2004). Nevertheless, we consider it likely that this estimate is accurate to within one order of magnitude, and that if it is not then it is more likely to be an underestimate than an overestimate. Based upon this we will consider in this paper the range of likely values to be $0.04 \leq A_{\text{trans,BHC}} \leq 4.0$.

For the transient sources we further need to consider periods in 'soft' X-ray states when the radio emission, and hence presumably the jet, are suppressed or 'quenched' (Fender et al. 1999; Corbel et al. 2001, 2004; Fender, Belloni & Gallo 2004; Fender 2005). The physical processes responsible for the suppression of the jet are not clear, but they seem to reduce the radio emission by a factor ≥ 30 – as a result we shall consider that when in such states the jet power drops to zero. The duty cycle in such states varies from outburst to outburst and it is hard to estimate a reasonable mean value. Based upon outbursts of the black hole LMXBs GX 339-4, XTE J1550-564 and XTE J1859+226 it seems a value of 60-80% for the duty cycle in 'soft' jet-suppressed states is reasonable (J. Homan, private communication; see also Fender, Belloni & Gallo 2004; Homan & Belloni 2005). As a result, we multiply the integrated power from transients by a factor 0.3. While this number is rather uncertain, we note that it is still the uncertainty in the value of the normalisation A_{trans} which dominates the integrated power input.

In the absence of sufficient good data on neutron star transient outburst with which to make a fit to $A_{\text{trans,NS}}$, we simply follow the approach adopted for the steady jets, i.e. $A_{\text{trans,NS}} = 0.1 A_{\text{trans,BHC}}$.

3 TOTAL POWER

As noted above, $P_J(L_X)$ is probably a function both of X-ray state and of the nature of the accreting compact star (neutron star or black hole). Convolved with both HMXB and LMXB luminosity functions, this presents us with eight populations of jet producing objects to consider (LMXB/HMXB $\times$ steady/transient $\times$ NS/BHC).

The integrated total jet power for any one of these classes is given by

$$L_{J,\text{total}} = f \int_{L_{X,\text{min}}}^{L_{X,\text{max}}} P_J \frac{dN}{dL_X} dL_X \quad (10)$$

where $P_J(L_X)$ is the jet power at a given X-ray luminosity L_X , and $N_X dL_X$ is the differential luminosity function for the class of object. The term f is a correction factor which can be separated into two parts:

$$f = f_{\text{pop}} f_{\text{spec}} \quad (11)$$

The first term, f_{pop} is simply the fraction of the population of that class of objects, in the sense that $f_{\text{pop,NS}} + f_{\text{pop,BHC}} = 1$. The second term, f_{spec} compensates for the fact that the X-ray luminosity functions given above, from Grimm et al. (2003) and Gilfanov (2004) consider the X-ray luminosity in the range ≤ 10 keV, whereas the expressions for the jet power were arrived at considering the total X-ray power. This term is therefore in some sense a bolometric correction. The values of f_{spec} for black holes are estimated in Portegies Zwart, Dewi & Maccarone (2004) for the black hole candidates. For the neutron star sources we followed essentially the same method. Estimated values for f_{pop} and f_{spec} are presented in table 1.

It is already clear from Fig 1 that HMXBs are unlikely to contribute significantly to the total power output, kinetic or radiative, of X-ray binaries (with the notable exceptions of SS 433 and Cyg X-3 which are powerful jet sources; see section 4). Simple calculations bear this out and from this point on we do not consider HMXBs further.

Radiative and kinetic output of LMXBs

In Fig 2 we present the integrated X-ray and jet (kinetic) luminosities of LMXBs as a function of source X-ray luminosity, scaled for the Milky Way, for three sets of values of jet power normalisations A_{steady} and A_{trans} . The shape of the functions result from the multiplication of the LMXB XLF (equation (1)) and the switch from steady to transient jets, which we fix to occur at 1.9×10^{37} for the black holes and 6×10^{36} for the neutron stars, e.g. around 20% of the Eddington limit for each class of object. Some features are common between the figures, and indeed independent of the selected values of the jet normalisation, namely:

- The radiative (X-ray) output is dominated by the more numerous NS LMXBs
- The kinetic (jet) output is dominated by the BHC LMXBs, which produce more jet power at a given L_X


The integrated bolometric radiative (X-ray) output of the two classes of objects is $L_{X,\text{bol,NS}} = 6.4 \times 10^{39} \text{ erg s}^{-1}$ and $L_{X,\text{bol,BHC}} = 1.7 \times 10^{39} \text{ erg s}^{-1}$.

Moving on to individual figures, Fig 2(a) ('case A') considers the 'minimum' or conservative estimates of the jet normalisation. In this scenario both classes of object, NS and BHC, produce more integrated radiative than kinetic output. Even with these low values for the normalisation, a transition from 'X-ray dominated' to 'Jet dominated' is apparent for the BHCs (see Fender, Gallo & Jonker 2003; Malzac, Merloni & Fabian 2004 for a fuller discussions). The integrated BHC jet luminosity is $L_J = 1.5 \times 10^{38} \text{ erg s}^{-1}$ (22% from transient jets).

Fig 2(b) ('case B') considers the situation where the steady jets reach power equipartition with the X-rays at the state transition, and dominate at lower X-ray luminosities (see e.g. Livio, Pringle & King 2003 for support for this point of view). The transient jets have a normalisation corresponding to the best fit from Fender, Belloni & Gallo (2004). In this scenario the peak of the kinetic power output from the BHC LMXBs is comparable to the peak of the radiative power output from the NS LMXBs. The integrated jet luminosity is $L_J = 2.0 \times 10^{39} \text{ erg s}^{-1}$ (19% from transient jets).

Table 1. Estimated values for the population f_{pop} and spectral f_{spec} corrections to be made to the integrals for the different classes of X-ray binary, as well as the jet power normalisation A . See text for more details.

		steady jets			transient jets		
		f_{pop}	f_{spec}	A	f_{pop}	f_{spec}	A
LMXB	NS	0.85	5.0	0.0006–0.03	0.85	2.0	0.0004–0.4
	BHC	0.15	10.5	0.006–0.3	0.15	1.2	0.04–4.0
HMXB	NS	0.95	5.0	0.0006–0.03	0.95	2.0	0.0
	BHC	0.05	10.0	0.006–0.3	0.05	1.2	0.04–4.0

**Figure 2.** The summed power output of all the Milky Way X-ray binaries, in radiation (X-rays) and kinetic power (jets), for three different sets of jet power normalisations, as a function of soft X-ray luminosity. The neutron stars dominate the radiative output but the kinetic power output is dominated by the black holes.**Table 2.** Supernova rates in SNU for different galaxy types

Galaxy type	Ia	Ib/c and II	Total
E-S0	0.18 ± 0.06	< 0.03	0.18 ± 0.06
S0a-Sb	0.18 ± 0.07	0.53 ± 0.19	0.72 ± 0.21
Sbc-Sd	0.21 ± 0.08	1.00 ± 0.35	1.21 ± 0.37

Fig 2(c) ('case C') is perhaps the most 'extreme' case, in which the kinetic output from the BHC LMXBs dominates the radiative output at essentially all L_X . The integrated jet luminosity is $L_J = 1.0 \times 10^{40} \text{ erg s}^{-1}$ (26% from transient jets).

4 JETS VS. SUPERNOVAE

A 'standard' type-I or type-II supernova explosion is believed to deposit around 10^{51} erg of kinetic energy into the surrounding interstellar medium (e.g. Chevalier 1977; Korpi et al. 1999). For an estimated rate of one such event per 100 years in the Milky Way, the time-averaged kinetic power is $L_{SNe} = 3 \times 10^{41} \text{ erg s}^{-1}$. Furthermore, Heinz & Sunyaev (2002) estimate the cosmic ray luminosity of the Milky Way as being approximately 10% of this, $L_{CR} = 4 \times 10^{40} \text{ erg s}^{-1}$.

Comparing these numbers with the three cases of jet power normalisation considered above, we can see that

$$10^{-3} \lesssim L_J/L_{SNe} \lesssim 0.03$$

$$10^{-2} \lesssim L_J/L_{CR} \lesssim 0.3$$

So it is clear that there may be enough power in the jets from X-ray binaries to supply a sizeable fraction of the cosmic ray luminosity of the Milky Way. There are many caveats to this, of course, but this number is of considerable interest. If we consider 'case B' outlined above to be the most likely, we find that the summed kinetic luminosity of X-ray binary jets is $\gtrsim 5\%$ of L_{CR} .

4.1 Early-type galaxies

So far we have only considered the situation for the Milky Way. However, different types of galaxies have different ratios of X-ray binaries to supernovae, which will affect the ratios calculated above. Estimated supernova rates for different types of galaxies are given in table 2 (from Cappellaro, Evans & Tuatto 1999).

Note that the total SNe rate in elliptical galaxies is about a factor of six less than in late-type spirals such as the Milky Way (due to the complete absence of SNe types Ib/c and II – see e.g. Cappellaro et al. 1999; van den Bergh, Li & Fillipenko 2002).

What do we expect of the X-ray binary, jet-producing,

population? Gilfanov (2004) has shown that the population of LMXBs scales linearly with the stellar mass of the galaxy, whereas for early-type galaxies the HMXB population is expected to be essentially zero. We should note that at least two of the most powerful jet sources, Cyg X-3 and SS 433, appear to be HMXBs (but in fact were not included in the preceding analysis as all HMXB contributions were disregarded). Nevertheless, GRS 1915+105 is a LMXB as are the seven most luminous neutron star XRB jet sources (the Z sources plus Cir X-1), as are all the BHC transients. Based on this we may crudely estimate that the jet power from 'outbursting' systems, per unit mass, is reduced by about a factor of no more than two in ellipticals compared to late-type spirals.

As an example we may take the case of NGC 4472, a large early-type galaxy. Gilfanov (2004) gives its stellar mass (the component which is correlated with the number of LMXBs) as being 3.5 times that of the Milky Way. Considering the scaling of LMXB numbers with stellar mass, as well as the possible 50% reduction in the brightest sources due to the lack of 'odd' HMXBs such as SS 433 and Cyg X-3, we would expect the summed jet power in NGC 4472 to be about twice that in the Milky Way. The total number of supernovae should also scale by mass, but is reduced by a factor ~ 6 due to the lack of any type Ib/c or type II events, and so would be around half of that in the Milky Way. Therefore the ratio of L_J/L_{SNe} would be ~ 4 times greater in this system, which may be considered to be typical for early-type ellipticals. In this situation, 'case C' outlined above would correspond to a total jet power input which was $\gtrsim 15\%$ of the power output of SNe. Even in the more conservative 'case B' the X-ray binary jets would be injecting a significant amount of power into the ISM.

CONCLUSIONS AND CAVEATS

In this paper we have demonstrated that the integrated kinetic power from X-ray binary jets in our galaxy is sufficient for them to be a significant source of cosmic rays. Observations have already clearly established that jets from powerful black hole transient XRBs can accelerate electrons to very high (TeV) energies (e.g. Corbel et al. 2002), but there remains little direct evidence for the acceleration of protons or atomic nuclei to relativistic energies (although heating of atomic nuclei to $\sim 10^7$ K at multiple sites in the jets of SS 433 has been demonstrated e.g. Marshall, Canizares & Schulz 2001 and references therein; Migliari, Fender & Mendez 2002). However, unequivocal evidence for acceleration of the non-leptonic component of cosmic rays in supernovae remnant shocks was also very thin on the ground until recently (Aharonian et al. 2004), and historically the argument was based upon energetics alone.

Given that the underlying physics of particle acceleration in SNR shocks and XRB jets / jet-ISM interactions is likely to be similar, if one process can accelerate protons to very high energies then presumably so can the other. It should furthermore be noted that XRB jets almost certainly spend a considerably longer phase moving with bulk relativistic velocities than do SNR, and the variation of efficiency of shock acceleration with bulk Lorentz factor is not well known. So it returns, at this stage, to simply a ques-

tion of energy budgets. Within the Milky Way, we conclude that the integrated jet power is likely to be around 1% of the rate of injection of energy from supernovae, which is around 5–10% of the cosmic ray luminosity, comparable to the numbers estimated by Heinz & Sunyaev (2002). There exists a distinct possibility that these numbers are serious underestimates. In addition, the total number of black holes in our galaxy accreting at very low levels remains highly uncertain, and the contribution from such sources is difficult to quantify; this is an area which needs some consideration.

The situation in the early-type galaxies, as illustrated with the example of NGC 4472, is therefore even more striking. In this system, with a mass 3.5 times that of the Milky Way, the total supernovae rate is likely to be only half that of our galaxy, yet with a population of LMXBs which scales as the mass. Even if the energisation of cosmic rays in our own galaxy is dominated by SNe, in such systems a significant fraction of the cosmic ray energy input should come from X-ray binary jets.

Furthermore, by assuming, as suggested by Gilfanov (2004), that the normalisation of the LMXB X-ray luminosity function is proportional to a galaxy's stellar mass, we have most likely underestimated the importance of LMXBs in giant elliptical galaxies and especially in the central dominant (cD) galaxies of clusters; some scatter is found in the relation of Gilfanov (2004), and in fact, the residuals are correlated with the globular cluster specific frequencies of the galaxies (White, Sarazin & Kulkarni 2002). Furthermore, it is known that globular cluster specific frequencies are generally higher in the most massive galaxies (e.g. McLaughlin, Harris & Hanes 1993), and the metal rich globular clusters, which are more likely to have X-ray binaries (Kundu, Maccarone & Zepf 2002) are even more prevalent in these systems (e.g. Kundu & Whitmore 2001) – the fraction of the X-ray sources in globular clusters clearly increases as one moves from S0 to giant elliptical to cD galaxies (Maccarone, Kundu & Zepf 2003). As a result, the most massive galaxies are likely to have somewhat higher L_{LMXB}/L_{NIR} ratios than the Milky Way, making the estimates here conservative.

There are of course many caveats and unknowns associated with the entirety of the above discussion, as there are with the assumption of SNe-dominated energisation. These include the energetic contribution from gamma-ray bursts and 'hypernovae' (e.g. Iwamoto et al. 1998), although the rates for such events are likely to be orders of magnitude less than those of core collapse supernovae (Podsiadlowski et al. 2004). Furthermore, there exists a distinct possibility that there is a poorly-known population of radiatively-faint sources with powerful jets such as LS 5039 (Paredes et al. 2000).

To conclude, we argue that the input of kinetic energy into the ISM in our galaxy may have a significant contribution from the jets of X-ray binaries, although is still likely to be dominated by supernovae. In early-type galaxies, which have no core collapse supernovae but a comparable number of low-mass X-ray binaries per unit stellar mass, the relative contribution will be much larger and should be considered as potentially rivalling supernovae. Given the high Lorentz factors of the most powerful X-ray binary jets, their effect may be most evident in its contribution to the production of cosmic rays. Finally, the energetic contribution of these

sources to the ISM will be in the form of a much more distributed source than the rare, more energetic, supernovae, which may have implications for the distribution of cosmic rays throughout the galaxy, and be an important input for models of their propagation through the galaxy (e.g. Strong, Moskalenko & Reimer 2004).

ACKNOWLEDGEMENTS

RPF would like to thank Marat Gilfanov, Hans-Jakob Grimm, Christian Kaiser, John Kirk and Ralph Wijers for useful conversations. Equally, TJM would like to thank Tiziana di Salvo, Chris Belczynski and Vicki Kalogera. RPF and TJM would both like to thank Jeoren Homan for a very useful contribution to our understanding of the typical duty cycles of different states in X-ray outbursts.

REFERENCES

- Aharonian F.A. et al., 2004, *Nature*, 432, 75
 Blandford R.D., Königl A., 1979, *ApJ*, 232, 34
 Cappellaro E., Evans R., Turatto M., 1999, *A&A* 351, 459
 Chevalier R., 1977, *ARA&A*, 15, 175
 Corbel S., Fender R.P., Tzioumis A.K., Nowak M., McIntyre V., Durouchoux P., Sood R., 2000, *A&A*, 359, 251
 Corbel S. et al., 2001, *ApJ*, 554, 43
 Corbel S., Nowak M., Fender R.P., Tzioumis A.K., Markoff S., 2003, *A&A*, 400, 1007
 Corbel S., Fender R.P., Tomsick J.A., Tzioumis A.K., Tingay S., 2004, *ApJ*, 617, 1272
 Elmegreen B.G., Scalo J., 2004, *ARA&A*, 42, 211
 Falcke H., Biermann P.L., 1996, *A&A*, 308, 321
 Fender R.P., 2001, *MNRAS*, 322, 31
 Fender R.P., 2005, in 'Compact Stellar X-Ray Sources', eds. W.H.G. Lewin and M. van der Klis, Cambridge University Press, (**astro-ph/0303339**)
 Fender R.P., Kuulkers E., 2001, *MNRAS*, 324, 923
 Fender R.P., Gallo E., Jonker P.G., 2003, *MNRAS*, 343, L99
 Fender R.P., Belloni T., Gallo E., 2004, *MNRAS*, 355, 1105
 Fender R.P. et al., 1999, *ApJ*, 519, L165
 Gallo E., Fender R.P., Pooley G.G., 2003, *MNRAS*, 344, 60
 Gilfanov M., 2004, *MNRAS*, 349, 146
 Grimm H.-J., Gilfanov M., Sunyaev R., 2002, *A&A*, 391, 923
 Grimm H.-J., Gilfanov M., Sunyaev R., 2003, *MNRAS*, 339, 793
 Heinz S., Sunyaev R., 2002, *A&A*, 390, 751
 Heinz S., Sunyaev R., 2003, *MNRAS*, 343, L59
 Homan J., Belloni T., to appear in *From X-ray Binaries to Quasars: Black Hole Accretion on All Mass Scales*, ed. T. J. Maccarone, R. P. Fender, L. C. Ho (Dordrecht: Kluwer). (**astro-ph/0412597**)
 Iwamoto K. et al., 1998, *Nature*, 395, 672
 Korpi M.J., Brandenburg A., Shukurov A., Tuominen I., Nordlund A., 1999, *ApJ*, 514, L99
 Kundu, A. & Whitmore, B.C., 2001, *AJ*, 121, 2950
 Kundu, A., Maccarone, T.J. & Zepf, S.E., 2002, *ApJL*, 574, 5
 Liu, Q. Z., van Paradijs J., van den Heuvel E. P. J., 2000, *A&AS*, 147, 25
 Liu, Q. Z., van Paradijs J., van den Heuvel E. P. J., 2001, *A&A*, 368, 1021
 McClintock J.E., Remillard R.A., 2005, in 'Compact Stellar X-Ray Sources', eds. W.H.G. Lewin and M. van der Klis, Cambridge University Press, (**astro-ph/0306213**)
 McLaughlin, D.E., Harris, W.E. & Hanes, D.A., 1993, *ApJL*, 409, 45
 Maccarone, T.J., Kundu, A., & Zepf, S.E., 2003, *ApJ*, 586, 814
 McCray R., Snow T.P., 1979, *ARA&A*, 17, 213
 Malzac J., Merloni A., Fabian A.C., 2004, *MNRAS*, 351, 253
 Markoff S., Falcke H., Fender R.P., 2001, *A&A*, 372, L25
 Marshall H.L., Canizares C.R., Schulz N.S., 2002, *ApJ*, 564, 941
 Migliari M., Fender R., Mendez M., 2002, *Science*, 297, 1673
 Migliari S., Fender R.P., Rupen M., Jonker P.G., Klein-Wolt M., Hjellming R.M., van der Klis M., 2003, 342, L67
 Mirabel, I.F., Rodríguez, L.F., 1999, *ARA&A*, 37, 409
 Muno M.P., Belloni T., Dhawan V., Mogran E.H., Remillard R.A., Rupen M.P., *ApJ*, submitted (**astro-ph/0411313**)
 Paredes J.M., Marti J., Ribo M., Massi M., 2000, *Science*, 288, 234
 Podislawski P., Mazzali P.A., Nomoto K., Lazzati D., Cappellaro E., 2004, *ApJ*, 607, L17
 Portegies Zwart, S.F., Dewi, J. & Maccarone, T., 2004, *MNRAS*, 355, 413
 Strong A.W., Moskalenko I.V., Reimer O., 2004, *ApJ*, 613, 962
 van den Bergh S., Li. W., Fillipenko A.V., 2002, *PASP*, 114, 820
 White, R.E. III, Sarazin C.L. & Kulkarni, S.R., 2002, *ApJL*, 571, 23